



## LOUTH COUNTY COUNCIL

### APPROPRIATE ASSESSMENT SCREENING REPORT

FOR

### PROPOSED PLAYGROUND, ARDEE ROAD, COLLON, CO. LOUTH

### VOLUME II. APPENDICES

24<sup>th</sup> July 2023

#### DOCUMENT ISSUE STATUS

REPORT ISSUE	REFERENCE NO.	DATE		
<b>FINAL</b>	409-01	24/07/2023		
TITLE	NAME	POSITION	SIGNATURE	DATE
<b>AUTHOR</b>	Leanne Tuohy	Staff Ecologist		18/07/2023
<b>MANAGING DIRECTOR</b>	Padraic Mulroy	Project Director		24/07/2023



## LIST OF APPENDICES


APP. No.	DESCRIPTION
1	Desk Study Information on Topsoils, Subsoils, Geology, Hydrogeology, Hydrology, Borehole Drilling Data and Historical Data from EPA, OSI, <a href="http://www.catchment.ie">www.catchment.ie</a> .
2	<i>NPWS (2021) Conservation Objectives: River Boyne and River Blackwater SAC 002299. Version 1. National Parks and Wildlife Service, Department of Housing, Local Government and Heritage.</i>  <i>NPWS (2022) Conservation objectives for River Boyne and River Blackwater SPA [004232]. First Order Site-specific Conservation Objectives Version 1.0. Department of Housing, Local Government and Heritage</i>



## **APPENDIX 1**

DESK STUDY INFORMATION ON TOPSOILS, SUBSOILS,  
GEOLOGY, HYDROGEOLOGY, HYDROLOGY, BOREHOLE  
DRILLING DATA & HISTORICAL DATA GATHERED FROM  
OSI, EPA, GSI & WWW.CATCHMENT.IE





Address/Eircode Search




3km

702,073.029 774,652.576 Meters



▼ Address/Eircode Search 🔍



60m

700,016.199 782,150.434 Meters





Crockanallis

4310  
413  
ST. S.  
ST. S.  
386

STREET  
MKT. SQ.  
School

Cath. Ch.

COL

Toberdillen

B.M. 407.1

400

Church  
Grave  
Yd.

B.M. 401.5

407

DROGHEDA

408 G.S.S.

School  
(Infants)  
School

400

60m



Address/Eircode Search



# COLLON

MARKET SQUARE  
W.M. 421

Court House

School  
B.M. 423.9

R.C. Church

Taberdillen  
27.868

Dispy. DROGHEDA

Constabulary Barrack

Church

Orange Hall

Rectory

School

60m

699,786.011 781,824.996 Meters



Crockanallis

4310  
413  
ST. S.  
ST. S.  
386

STREET  
MKT. SQ.  
School

Cath. Ch.

COL

Toberdillen

B.M. 407.1

400

Church  
Grave  
Yd.

B.M. 401.5

DROGHEDA

408 G.S.S.

School  
(Infants)  
School

400

60m



# Results

[Link to More Information](#)



## SIS National Soils

### Urban

Association_Name	Urban
Association_Unit	Urban
Association_Symbol	Urban
Texture_Substrate_Type	Urban
Ha	13.29468059
Drainage	Other
Texture	
Depth	
SOC	
URL	<a href="#">Link to More Information</a>



EXPORT



 Keep Previous Results**SIS National Soils  
Clonroche**

Association_Name	Clonroche
Association_Unit	1100a
Association_Symbol	1100a
Texture_Substrate_Type	Fine loamy drift with sillaceous stones
Ha	13459.64623268
Drainage	Well
Texture	Fine loamy
Depth	>80
SOC	129.44465676
URL	<a href="#">Link to More Information</a>

**SIS National Soils  
Urban**

1



2 km




# Results

URL [Link to More Information](#)

SIS National Soils River	
Association_Name	River
Association_Unit	05RIV
Association_Symbol	05RIV
Texture_Substrate_Type	River alluvium
Ha	47.49964894
Drainage	Poor
Texture	All
Depth	>80
SOC	173.76511738
URL	<a href="#">Link to More Information</a>

EXPORT





Quaternary Sediments 1:50,000 Ireland ITM

Lithology	Till derived from Lower Palaeozoic sandstones and shales
Quaternary Sediment	TLPSsS

[Zoom to](#) ⋮



# Results



Keep Previous Results

## National Soils Hydrology Map Poorly Drained



CATEGORY	Poorly Drained
SOILDRAINAGECLASS	Poor
PAR_MAT	TLPSsS
IFS_SOIL	AminPD
IFS_CODE	31



## National Soils Hydrology Map Made



## National Soils Hydrology Map Poorly Drained



EXPORT



1



OSi




HELP



500 m





(1 of 2) ▶ □ ✕


**Quaternary Sediments 1:50,000 Ireland ITM**

---

Lithology

- Alluvium
- Quaternary Sediment A

[Zoom to](#) ...



Quaternary Sediments 1:50,000 Ireland ITM


Lithology	Gravels derived from Lower Palaeozoic sandstones and shales
Quaternary Sediment	<a href="#">GLPSsS</a>

[Zoom to](#) ⋮

0.6km

Map navigation and search controls:

- Zoom in (+) and zoom out (-) buttons.
- Search bar: Enter Address, Eircode or T
- Home button (house icon)
- Refresh button (circular arrow icon)
- Measurement tool (ruler icon)
- Mobile view button (phone icon)
- Layers button (layers icon)
- Full screen button (expand icon)



(1 of 2) [Navigation icons]

**Quaternary Sediments 1:50,000 Ireland ITM**

Lithology	Bedrock outcrop or subcrop
Quaternary Sediment	Rck

[Zoom to](#) [More options icon]

 Keep Previous Results

GSI Bedrock Polygons 100k

**Collon Formation**

UnitName	Collon Formation
NewCode	COLN
SheetNumber	13
StratigraphicCode	CM
LithologicalCode	
Description	Andesite breccia/conglomerate/sandstone
Label	CM
AreaKm2	1.07053005
PerimeterM	4633.10009765



EXPORT



1



GSI



HELP



500 m



# Results



5



Keep Previous Results

## National Water Monitoring Stations GWIE\_EA\_G\_01021000004



StationID	GWIE_EA_G_01021000004
StationName	Collon
StationType	GROUNDWATER_STATION
WFDWISECODE	IEMGGWIE_EA_G_01021000004
EntityCode	IE_EA_G_010
EntityName	Wilkinstown
WBWFDWISECODE	IE_EA_G_010
TypeofWaterMonitored	Groundwater body code but there is no corresponding code in Eden MDS for the moment
LocalAuthority	LOUTH COUNTY COUNCIL
RiverBasinDistrict	Eastern
EPASStationTypeWFDs	SurveillanceAndOperational
CreatedByOrganisation	EPA
EPALink	N/A
Easting	300239
Northing	281687.28

OSi



100 m



Results



1



HELP



Keep Previous Results

GSI Rock Unit Groups

Ordovician Volcanics



NAME

Ordovician Volcanics

CalcClass

Non-Calcareous



EXPORT

OSi

500 m



Results



1



HELP



Keep Previous Results

**GSI Bedrock Geology 1 Million  
Ordovician volcanic rocks**



UnitName	Ordovician volcanic rocks
GeologicalAge	Palaeozoic, Ordovician
AreaKm2	2.40757915
Perimeter	8111.33984375



EXPORT

GSI

500 m





Results



Keep Previous Results

GSI Bedrock Aquifer

Pl

AquiferCode Pl

AquiferDesc Poor Aquifer - Bedrock which is Generally Unproductive except for Local Zones



EXPORT



HELP



GSI

500 m



Results



Keep Previous Results

GSI Bedrock Aquifer

Pu

AquiferCode

Pu

AquiferDesc

Poor Aquifer - Bedrock which is Generally Unproductive

EXPORT



HELP



GSI

500 m



# Results



Keep Previous Results

GSI Bedrock Aquifer



**Pu**

AquiferCode


Pu

AquiferDesc

Poor Aquifer - Bedrock which is Generally Unproductive



EXPORT





# Results



Keep Previous Results

## WFD SubCatchments




### Boyne\_SC\_120

Name	Boyne_SC_120
Subcatchment_Id	07_15
Catchment_Id	07
CreatedOn	2015-08-23T23:00:00Z
UpdatedOn	2017-03-07T00:00:00Z
Local_Authority	MEATH COUNTY COUNCIL



EXPORT



# Results



Keep Previous Results


## WFD River Sub Basins MATTOCK\_010



NAME	MATTOCK_010
EU_CD	IE_EA_07M010100
AREAKM2	16.64034996




EXPORT



Results

Keep Previous Results

River Waterbodies	
IE_EA_07M010100	
EU_CD	IE_EA_07M010100
NAME	MATTOCK_010
URL	<a href="#">View the Data Page</a>
MS_CD	EA_07M010100
REGION_CD	17
INS_WHEN	2014-11-14T00:00:00Z
BASIN_CD	159 Boyne
LAT	53.76765268
LON	-6.50671191
LENGTHKM	18.09094885
SUB_CD	07_15
DateChanged	2019-11-20T00:00:00Z
Change	Rivers updated
GEOLOGY	1
CatchmentAreaKm2	16.64034996
CatchmentAreaHectares	1664.034996
Slope	-0.00773877




HELP



Results

Latest River Q Values  
**RS07M010050**


StationCode	RS07M010050
StationName	MATTOCK - Slane Rd Br Collon
StationTypeEDEN	RIVER_STATION
RiverWaterbodyName	MATTOCK_010
EntityName	MATTOCK
EntityCode	07M01
Year	2020
QValueScore	4
QValueStatus	Good
WFDWISECODE	IEMRRS07M010050
WBWFDWISECODE	IE_EA_07M010100
LocalAuthority	LOUTH COUNTY COUNCIL
EPASStationTypeWFD	Operational
Typeofwatermonitored	River Water
RiverBasinDistrict	Eastern
SegCd	07_1405





Results

Latest River Q Values	
RS07M010050	
StationCode	RS07M010050
StationName	MATTOCK - Slane Rd Br Collon
StationTypeEDEN	RIVER_STATION
RiverWaterbodyName	MATTOCK_010
EntityName	MATTOCK
EntityCode	07M01
Year	2020
QValueScore	4
QValueStatus	Good
WFDWISECODE	IEMRRS07M010050
WBWFDWISECODE	IE_EA_07M010100
LocalAuthority	LOUTH COUNTY COUNCIL
EPASStationTypeWFD	Operational
Typeofwatermonitored	River Water
RiverBasinDistrict	Eastern
SegCd	07_1405





## **APPENDIX 2**

- NPWS (2021) CONSERVATION OBJECTIVES: RIVER BOYNE AND RIVER BLACKWATER SAC 002299. VERSION 1.  
NATIONAL PARKS AND WILDLIFE SERVICE,  
DEPARTMENT OF HOUSING, LOCAL GOVERNMENT  
AND HERITAGE.
- NPWS (2022) CONSERVATION OBJECTIVES FOR RIVER BOYNE AND RIVER BLACKWATER SPA [004232].  
FIRST ORDER SITE-SPECIFIC CONSERVATION  
OBJECTIVES VERSION 1.0.  
DEPARTMENT OF HOUSING, LOCAL GOVERNMENT  
AND HERITAGE



## **SITE SYNOPSIS**

**SITE NAME: RIVER BOYNE AND RIVER BLACKWATER SPA**

**SITE CODE: 004232**

The River Boyne and River Blackwater SPA is a long, linear site that comprises stretches of the River Boyne and several of its tributaries; most of the site is in Co. Meath, but it extends also into Cos Cavan, Louth and Westmeath. It includes the following river sections: the River Boyne from the M1 motorway bridge, west of Drogheda, to the junction with the Royal Canal, west of Longwood, Co Meath; the River Blackwater from its junction with the River Boyne in Navan to the junction with Lough Ramor in Co. Cavan; the Tremblestown River/Athboy River from the junction with the River Boyne at Kilnagross Bridge west of Trim to the bridge in Athboy, Co. Meath; the Stoneyford River from its junction with the River Boyne to Stonestown Bridge in Co. Westmeath; the River Deel from its junction with the River Boyne to Cummer Bridge, Co. Westmeath. The site includes the river channel and marginal vegetation.

Most of the site is underlain by Carboniferous limestone but Silurian quartzite also occurs in the vicinity of Kells and Carboniferous shales and sandstones close to Trim.

The site is a Special Protection Area (SPA) under the E.U. Birds Directive of special conservation interest for the following species: Kingfisher.

A survey in 2010 recorded 19 pairs of Kingfisher (based on 15 probable and 4 possible territories) in the River Boyne and River Blackwater SPA. A survey conducted in 2008 recorded 20-22 Kingfisher territories within the SPA. Other species which occur within the site include Mute Swan (90), Teal (166), Mallard (219), Cormorant (36), Grey Heron (44), Moorhen (84), Snipe (32) and Sand Martin (553) – all figures are peak counts recorded during the 2010 survey.

The River Boyne and River Blackwater Special Protection Area is of high ornithological importance as it supports a nationally important population of Kingfisher, a species that is listed on Annex I of the E.U. Birds Directive.

25.11.2010



## **Conservation objectives for River Boyne and River Blackwater SPA [004232]**

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Objective: To maintain or restore the favourable conservation condition of the bird species listed as Special Conservation Interests for this SPA:

<b>Bird Code</b>	<b>Common Name</b>	<b>Scientific Name</b>
A229	Kingfisher	<i>Alcedo atthis</i>



**Citation:** NPWS (2022) *Conservation objectives for River Boyne and River Blackwater SPA [004232]. First Order Site-specific Conservation Objectives Version 1.0. Department of Housing, Local Government and Heritage.*

*This First Order Site-specific Conservation Objectives Version 1.0 document replaces the Generic Conservation Objectives Version 9.0 document.*







**Site Name: River Boyne and River Blackwater SAC**

**Site Code: 002299**

This site comprises the freshwater element of the River Boyne as far as the Boyne Aqueduct, the Blackwater as far as Lough Ramor and the Boyne tributaries including the Deel, Stoneyford and Tremblestown Rivers. These riverine stretches drain a considerable area of Meath and Westmeath, and smaller areas of Cavan and Louth. The underlying geology is Carboniferous Limestone for the most part, with areas of Upper, Lower and Middle well represented. In the vicinity of Kells Silurian Quartzite is present while close to Trim are Carboniferous Shales and Sandstones. There are many large towns adjacent to but not within the site, including Slane, Navan, Kells, Trim, Athboy and Ballivor.

The site is a Special Area of Conservation (SAC) selected for the following habitats and/or species listed on Annex I / II of the E.U. Habitats Directive (\* = priority; numbers in brackets are Natura 2000 codes):

- |  |
|--|
| [7230] Alkaline Fens                                 |
| [91E0] Alluvial Forests*                             |
| [1099] River Lamprey ( <i>Lampetra fluviatilis</i> ) |
| [1106] Atlantic Salmon ( <i>Salmo salar</i> )        |
| [1355] Otter ( <i>Lutra lutra</i> )                  |

The main areas of alkaline fen in this site are concentrated in the vicinity of Lough Shesk, Freehan Lough and Newtown Lough. The hummocky nature of the local terrain produces frequent springs and seepages which are rich in lime. A series of base-rich marshes have developed in the poorly-drained hollows, generally linked with these three lakes. Open water is usually fringed by Bulrush (*Typha latifolia*), Common Club-rush (*Scirpus lacustris*) or Common Reed (*Phragmites australis*), and this last species also extends shorewards where a dense stand of Great Fen-sedge (*Cladium mariscus*) frequently occurs. This in turn grades into a sedge and grass community (*Carex* spp. and Purple Moor-grass, *Molinia caerulea*), or one dominated by Black Bog-rush (*Schoenus nigricans*). An alternative aquatic/terrestrial transition is a floating layer of vegetation. This is normally based on Bogbean (*Menyanthes trifoliata*) and Marsh Cinquefoil (*Potentilla palustris*). Other species gradually become established on this cover, especially plants tolerant of low nutrient status e.g. bog mosses (*Sphagnum* spp.). Diversity of plant and animal life is high in the fen and the flora includes many rarities. Plants of interest include Narrow-leaved Marsh-orchid (*Dactylorhiza traunsteineri*), Fen Bedstraw (*Galium uliginosum*), Cowbane (*Cicuta virosa*), Frogbit (*Hydrocharis morsus-ranae*) and Least Bur-reed (*Sparganium minimum*). These species tend to be restricted in their distribution in Ireland. Also notable is the

abundance of aquatic stoneworts (*Chara* spp.) which are characteristic of calcareous wetlands.

The rare plant Round-leaved Wintergreen (*Pyrola rotundifolia*) occurs around Newtown Lough. This species is listed in the Red Data Book and this site represents its only occurrence in Co. Meath.

Wet woodland fringes many stretches of the Boyne. The Boyne River Islands are a small chain of three islands situated 2.5 km west of Drogheda. The islands were formed by the build-up of alluvial sediment in this part of the river where water movement is sluggish. All of the islands are covered by dense thickets of wet, willow (*Salix* spp.) woodland, with the following species occurring: Osier (*S. viminalis*), Crack Willow (*S. fragilis*), White Willow (*S. alba*), Purple Willow (*Salix purpurea*) and Rusty Willow (*S. cinerea* subsp. *oleifolia*). A small area of Alder (*Alnus glutinosa*) woodland is found on soft ground at the edge of the canal in the north-western section of the islands. Along other stretches of the rivers of the site Rusty Willow scrub and pockets of wet woodland dominated by Alder have become established, particularly at the river edge of mature deciduous woodland. Ash (*Fraxinus excelsior*) and Downy Birch (*Betula pubescens*) are common in the latter, and the ground flora is typical of wet woodland with Meadowsweet (*Filipendula ulmaria*), Wild Angelica (*Angelica sylvestris*), Yellow Iris (*Iris pseudacorus*), horsetails (*Equisetum* spp.) and occasional tussocks of Greater Tussock-sedge (*Carex paniculata*).

The dominant habitat along the edges of the river is freshwater marsh, and the following plant species occur commonly in these areas: Yellow Iris, Creeping Bent (*Agrostis stolonifera*), Canary Reed-grass (*Phalaris arundinacea*), Marsh Bedstraw (*Galium palustre*), Water Mint (*Mentha aquatica*) and Water Forget-me-not (*Myosotis scorpioides*). In the wetter areas Common Meadow-rue (*Thalictrum flavum*) is found. In the vicinity of Dowth, Fen Bedstraw (*Galium uliginosum*), a scarce species mainly confined to marshy areas in the midlands, is common in this vegetation. Swamp Meadow-grass (*Poa palustris*) is an introduced plant which has spread into the wild (naturalised) along the Boyne approximately 5 km south-west of Slane. It is a rare species which is listed in the Red Data Book and has been recorded among freshwater marsh vegetation on the banks of the Boyne in this site. The only other record for this species in the Republic of Ireland is from a site in Co. Monaghan.

The secondary habitat associated with the marsh is wet grassland and species such as Tall Fescue (*Festuca arundinacea*), Silverweed (*Potentilla anserina*), Creeping Buttercup (*Ranunculus repens*), Meadowsweet and Meadow Vetchling (*Lathyrus pratensis*) are well represented. Strawberry Clover (*Trifolium fragiferum*), a plant generally restricted to coastal locations in Ireland, has been recorded from wet grassland vegetation at Trim. At Rosnaree river bank on the River Boyne, Round-Fruited Rush (*Juncus compressus*) is found in alluvial pasture, which is generally periodically flooded during the winter months. This rare plant is only found in three counties in Ireland.

Along much of the Boyne and along tributary stretches are found areas of mature deciduous woodland on the steeper slopes above the floodplain marsh or wet woodland vegetation. Many of these are planted in origin. However the steeper areas of King Williams Glen and Townley Hall wood have been left unmanaged and now have a more natural character. East of Curley Hole the woodland has a natural appearance with few conifers. Broadleaved species include oaks (*Quercus* spp.), Ash, willows, Hazel (*Corylus avellana*), Sycamore (*Acer pseudoplatanus*), Holly (*Ilex aquifolium*), Horse-chestnut (*Aesculus hippocastanum*) and the shrubs Hawthorn (*Crataegus monogyna*), Blackthorn (*Prunus spinosa*) and Elder (*Sambucus nigra*). South-west of Slane and in Dowth, some more exotic tree species such as Beech (*Fagus sylvatica*), and occasionally Lime (*Tilia cordata*), are seen. The coniferous trees Larch (*Larix* sp.) and Scots Pine (*Pinus sylvestris*) also occur. The woodland ground flora includes Barren Strawberry (*Potentilla sterilis*), Enchanter's-nightshade (*Circaea lutetiana*) and Ground-ivy (*Glechoma hederacea*), along with a range of ferns. Variation occurs in the composition of the canopy - for example, in wet patches alongside the river, White Willow and Alder form the canopy.

Other habitats present along the Boyne and Blackwater include lowland dry grassland, improved grassland, reedswamp, weedy waste ground, scrub, hedge, drainage ditch and canal. In the vicinity of Lough Shesk, the dry slopes of the morainic hummocks support grassland vegetation which, in some places, is partially colonised by Gorse (*Ulex europaeus*) scrub. Those grasslands which remain unimproved for pasture are species-rich, with Common Knapweed (*Centaurea nigra*), Creeping Thistle (*Cirsium arvense*) and Ribwort Plantain (*Plantago lanceolata*) commonly present. Fringing the canal alongside the Boyne south-west of Slane are areas with Reed Sweet-grass (*Glyceria maxima*), Great Willowherb (*Epilobium hirsutum*) and Meadowsweet.

The Boyne and its tributaries form one of Ireland's premier game fisheries and the area offers a wide range of angling, from fishing for spring salmon and grilse to seatrout fishing and extensive brown trout fishing. Atlantic Salmon (*Salmo salar*) use the tributaries and headwaters as spawning grounds. Although this species is still fished commercially in Ireland, it is considered to be endangered or locally threatened elsewhere in Europe and is listed on Annex II of the Habitats Directive. Atlantic Salmon run the Boyne almost every month of the year. The Boyne is most important as it represents an eastern river which holds large three-sea-winter fish from 20-30 lb. These fish generally arrive in February, with smaller spring fish (10 lb) arriving in April/May. The grilse come in July, water permitting. The river gets a further run of fish in late August and this run would appear to last well after the fishing season. The salmon fishing season lasts from 1<sup>st</sup> March to 30<sup>th</sup> September.

The Blackwater is a medium sized limestone river which is still recovering from the effects of the arterial drainage scheme of the 1970s. Salmon stocks have not recovered to the numbers that existed pre-drainage. The Deel, Riverstown, Stoneyford and Tremblestown Rivers are all spring-fed, with a continuous high volume of water. They are difficult to fish because some areas are overgrown, while others have been affected by drainage with resultant high banks.

This site is also important for the populations of two other species listed on Annex II of the E.U. Habitats Directive which it supports, namely River Lamprey (*Lampetra fluviatilis*), which is present in the lower reaches of the Boyne River, and Otter (*Lutra lutra*), which can be found throughout the site. In addition, the site also supports many more of the mammal species occurring in Ireland. Those which are listed in the Irish Red Data Book include Pine Marten, Badger and Irish Hare. Common Frog, another Red Data Book species, also occurs within the site. All of these animals, with the addition of the Stoat and Red Squirrel, which also occur within the site, are protected under the Wildlife Act, 1976.

Whooper Swans winter regularly at several locations along the Boyne and Blackwater Rivers. Known sites are at Newgrange (approx. 20 in recent winters), near Slane (20+ in recent winters), Wilkinstown (several records of 100+) and River Blackwater from Kells to Navan (104 at Kells in winter 1996/97, 182 at Headfort in winter 1997/98, 200-300 in winter 1999/00). The available information indicates that there is a regular wintering population of Whooper Swans based along the Boyne and Blackwater River valleys. The birds use a range of feeding sites but roosting sites are not well known. The population is substantial, certainly of national, and at times international, importance. Numbers are probably in the low hundreds.

Intensive agriculture is the main land use along the site. Much of the grassland is in very large fields and is improved. Silage harvesting is carried out. The spreading of slurry and fertiliser poses a threat to the water quality of this salmonid river and to the lakes. In the more extensive agricultural areas sheep grazing is carried out.

Fishing is a main tourist attraction on the Boyne and Blackwater and there are a number of Angler Associations, some with a number of beats. Fishing stands and styles have been erected in places. The Eastern Regional Fishery Board have erected fencing along selected stretches of the river as part of their salmonid enhancement programme. Parts of the river system have been arterially dredged. In 1969 an arterial dredging scheme commenced and disrupted angling for 18 years. The dredging altered the character of the river completely and resulted in many areas in very high banks. The main channel from Drogheda upstream to Navan was left untouched, as were a few stretches on the Blackwater. Ongoing maintenance dredging is carried out along stretches of the river system where the gradient is low. This is extremely destructive to salmonid habitat in the area. Drainage of the adjacent river systems also impacts on the many small wetland areas throughout the site. The River Boyne is a designated Salmonid Water under the E.U. Freshwater Fish Directive.

The site supports populations of several species listed on Annex II of the E.U. Habitats Directive, and habitats listed on Annex I of this Directive, as well as examples of other important habitat types. Although the wet woodland areas appear small there are few similar examples of this type of alluvial wet woodland remaining in the country, particularly in the north-east. The semi-natural habitats, particularly the strips of woodland which extend along the river banks, and the marsh and wet

grasslands, increase the overall habitat diversity and add to the ecological value of the site, as does the presence of a range of Red Data Book plant and animal species and the presence of nationally rare plant species.

# National Parks and Wildlife Service

---

## *Conservation Objectives Series*

---

### River Boyne and River Blackwater SAC 002299



An Roinn Tithíochta,  
Rialtais Áitiúil agus Oidhreachta  
Department of Housing,  
Local Government and Heritage

**National Parks and Wildlife Service,  
Department of Housing, Local Government and Heritage,  
90 King Street North, Dublin 7, D07 N7CV, Ireland.  
Web: [www.npws.ie](http://www.npws.ie)  
E-mail: [natureconservation@housing.gov.ie](mailto:natureconservation@housing.gov.ie)**

**Citation:**

**NPWS (2021) Conservation Objectives: River Boyne and River Blackwater SAC  
002299. Version 1. National Parks and Wildlife Service, Department of Housing,  
Local Government and Heritage.**

**Series Editors: Rebecca Jeffrey and Christina Campbell  
ISSN 2009-4086**

## Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

### Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.



## Qualifying Interests

\* indicates a priority habitat under the Habitats Directive

---

002299	River Boyne and River Blackwater SAC
1099	River Lamprey <i>Lampetra fluviatilis</i>
1106	Salmon <i>Salmo salar</i>
1355	Otter <i>Lutra lutra</i>
7230	Alkaline fens
91E0	Alluvial forests with <i>Alnus glutinosa</i> and <i>Fraxinus excelsior</i> (Alno-Padion, Alnion incanae, Salicion albae)*

**Please note that this SAC overlaps with Boyne Estuary SPA (004080) and River Boyne and River Blackwater SPA (004232). The SAC is also adjacent to Boyne Coast and Estuary SAC (001957). See map 2. The conservation objectives for this site should be used in conjunction with those for the overlapping and adjacent sites as appropriate.**

## Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: [www.npws.ie/Publications](http://www.npws.ie/Publications)

### NPWS Documents

<b>Year :</b>	2006
<b>Title :</b>	Otter survey of Ireland 2004/2005
<b>Author :</b>	Bailey, M.; Rochford, J.
<b>Series :</b>	Irish Wildlife Manuals, No. 23
<b>Year :</b>	2006
<b>Title :</b>	A baseline survey of juvenile lamprey populations in the Boyne catchment
<b>Author :</b>	O'Connor, W.
<b>Series :</b>	Irish Wildlife Manuals, No. 24
<b>Year :</b>	2007
<b>Title :</b>	Supporting documentation for the Habitats Directive Conservation Status Assessment - backing documents. Article 17 forms and supporting maps
<b>Author :</b>	NPWS
<b>Series :</b>	Unpublished report to NPWS
<b>Year :</b>	2008
<b>Title :</b>	National survey of native woodlands 2003-2008
<b>Author :</b>	Perrin, P.M.; Martin, J.; Barron, S.; O'Neill, F.H.; McNutt, K.E.; Delaney, A.
<b>Series :</b>	Unpublished report to NPWS
<b>Year :</b>	2009
<b>Title :</b>	Ireland Red List No. 2: Non-marine molluscs
<b>Author :</b>	Byrne, A.; Moorkens, E.A.; Anderson, R.; Killeen, I.J.; Regan, E.C.
<b>Series :</b>	Ireland Red List series, NPWS
<b>Year :</b>	2010
<b>Title :</b>	A provisional inventory of ancient and long-established woodland in Ireland
<b>Author :</b>	Perrin, P.M.; Daly, O.H.
<b>Series :</b>	Irish Wildlife Manuals, No. 46
<b>Year :</b>	2010
<b>Title :</b>	Ireland Red List No. 4: Butterflies
<b>Author :</b>	Regan, E.C.; Nelson, B.; Aldwell, B.; Bertrand, C.; Bond, K.; Harding, J.; Nash, D.; Nixon, D.; Wilson, C.J.
<b>Series :</b>	Ireland Red List series, NPWS
<b>Year :</b>	2012
<b>Title :</b>	Ireland Red List No. 8: Bryophytes
<b>Author :</b>	Lockhart, N.; Hodgetts, N.; Holyoak, D.
<b>Series :</b>	Ireland Red List series, NPWS
<b>Year :</b>	2013
<b>Title :</b>	National otter survey of Ireland 2010/12
<b>Author :</b>	Reid, N.; Hayden, B.; Lundy, M.G.; Pietravalle, S.; McDonald, R.A.; Montgomery, W.I.
<b>Series :</b>	Irish Wildlife Manuals, No. 76
<b>Year :</b>	2013
<b>Title :</b>	Results of a monitoring survey of old sessile oak woods and alluvial forests
<b>Author :</b>	O'Neill, F.H.; Barron, S.J.
<b>Series :</b>	Irish Wildlife Manuals, No. 71

**Year :** 2013  
**Title :** The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments  
**Author :** NPWS  
**Series :** Conservation assessments

---

**Year :** 2016  
**Title :** Ireland Red List No. 10: Vascular Plants  
**Author :** Wyse Jackson, M.; FitzPatrick, Ú.; Cole, E.; Jebb, M.; McFerran, D.; Sheehy Skeffington, M.; Wright, M.  
**Series :** Ireland Red Lists series, NPWS

---

**Year :** in prep.  
**Title :** The monitoring and assessment of four EU Habitats Directive Annex I woodland habitats  
**Author :** Daly, O.H.; O'Neill, F.H.; Barron, S.J.  
**Series :** Irish Wildlife Manuals

---

**Year :** in prep.  
**Title :** Scoping study and pilot survey of fens  
**Author :** O'Neill, F.H.; Perrin, P.M.; Denyer, J.; Martin, J.R.; Daly, O.H.; Brophy, J.T.  
**Series :** Irish Wildlife Manuals

## Other References

**Year :** 1982  
**Title :** Otter survey of Ireland  
**Author :** Chapman, P.J.; Chapman, L.L.  
**Series :** Unpublished report to Vincent Wildlife Trust

---

**Year :** 1991  
**Title :** The spatial organization of otters (*Lutra lutra*) in Shetland  
**Author :** Kruuk, H.; Moorhouse, A.  
**Series :** Journal of Zoology, 224: 41-57

---

**Year :** 2002  
**Title :** Reversing the habitat fragmentation of British woodlands  
**Author :** Peterken, G.  
**Series :** WWF-UK, London

---

**Year :** 2006  
**Title :** Otters - ecology, behaviour and conservation  
**Author :** Kruuk, H.  
**Series :** Oxford University Press

---

**Year :** 2010  
**Title :** Otter tracking study of Roaringwater Bay  
**Author :** De Jongh, A.; O'Neill, L.  
**Series :** Unpublished draft report to NPWS

---

**Year :** 2011  
**Title :** Comparison of field- and GIS-based assessments of barriers to Atlantic salmon migration: a case study in the Nore Catchment, Republic of Ireland  
**Author :** Gargan, P.G.; Roche, W.K.; Keane, S.; King, J.J.; Cullagh, A.; Mills, P.; O'Keeffe, J.  
**Series :** Journal of Applied Ichthyology, 27 (Suppl. 3): 66-72

**Year :** 2011  
**Title :** Review and revision of empirical critical loads and dose-response relationships. Proceedings of an expert workshop, Noordwijkerhout, 23-25 June 2010  
**Author :** Bobbink, R.; Hettelingh, J.P.  
**Series :** RIVM report 680359002, Coordination Centre for Effects, National Institute for Public Health and the Environment (RIVM)

---

**Year :** 2011  
**Title :** The Fen Management Handbook  
**Author :** McBride, A.; Diack, I.; Droy, N.; Hamill, B.; Jones, P.; Schutten, J.; Skinner, A.; Street, M. (eds.)  
**Series :** Scottish Natural Heritage, Perth

---

**Year :** 2015  
**Title :** Behaviour of sea lamprey (*Petromyzon marinus* L.) at man-made obstacles during upriver spawning migration: use of telemetry to access efficacy of weir modifications for improved passage  
**Author :** Rooney, S.M.; Wightman, G.D.; O Conchuir, R.; King, J.J.  
**Series :** Biology and Environment: Proceedings of the Royal Irish Academy, 115B: 1-12

---

**Year :** 2015  
**Title :** Common standards monitoring guidance for freshwater fauna. Version October 2015  
**Author :** JNCC  
**Series :** Joint Nature Conservation Committee, Peterborough

---

**Year :** 2016  
**Title :** Irish Vegetation Classification: Technical Progress Report No. 2  
**Author :** Perrin, P.  
**Series :** Report submitted to National Biodiversity Data Centre

---

**Year :** 2016  
**Title :** National Programme: Habitats Directive and Red Data Book Species Summary Report 2015  
**Author :** Gallagher, T.; O'Gorman, N.M.; Rooney, S.M.; Coghlan, B.; King, J.J.  
**Series :** IFI/2016/1-4344. Inland Fisheries Ireland

---

**Year :** 2018  
**Title :** Irish Vegetation Classification: Technical Progress Report No. 4  
**Author :** Perrin, P.  
**Series :** Report submitted to National Biodiversity Data Centre

---

**Year :** 2021  
**Title :** The Status of Irish Salmon Stocks in 2020 with Catch Advice for 2021  
**Author :** Gargan, P.; Fitzgerald, C.; Kennedy, R.; Maxwell, H.; McLean, S.; Millane, M.  
**Series :** Report of the Technical Expert Group on Salmon (TEGOS) to the North-South Standing Scientific Committee for Inland Fisheries

---

## Spatial data sources

**Year :** Revision 2010  
**Title :** National Survey of Native Woodlands 2003-2008. Version 1  
**GIS Operations :** QIs selected; clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising  
**Used For :** 91E0 (map 3)

---

**Year :** 2018  
**Title :** Woodland Monitoring Survey 2017-2018  
**GIS Operations :** QIs selected; clipped to SAC boundary. Expert opinion used as necessary to resolve any issues arising  
**Used For :** 91E0 (map 3)

---

## Conservation Objectives for : River Boyne and River Blackwater SAC [002299]

### 7230 Alkaline fens

**To maintain the favourable conservation condition of Alkaline fens in River Boyne and River Blackwater SAC, which is defined by the following list of attributes and targets:**

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Alkaline fen has not been mapped in detail for River Boyne and River Blackwater SAC and thus the exact total current area of the qualifying habitat in the SAC is currently unknown. The main areas of alkaline fen in the SAC are documented to occur in the vicinity of Lough Shesk, Freekan Lough, Newtown Lough in the upper reaches of the Stonyford River. At Lough Shesk, the habitat is particularly well-represented and there is a good example of succession from open water to fen-type habitat (NPWS internal files)
Habitat distribution	Occurrence	No decline, subject to natural processes	See the notes for habitat area above
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil pH and nutrient status within natural ranges	Relevant nutrients and their natural ranges are yet to be defined. However, nitrogen deposition is noted as being relevant to this habitat in NPWS (2013). See also Bobbink and Hettelingh (2011). Increased nutrients can lead to changes in plant and invertebrate species through competition and subsequent structural changes to micro-habitat. These nutrients favour growth of grasses rather than forbs and mosses and leads to a higher and denser sward
Ecosystem function: peat formation	Percentage cover of peat-forming vegetation and water table levels	Maintain active peat formation, where appropriate	In order for peat to form, water levels need to be slightly below or above the soil surface for c.90% of the time
Ecosystem function: hydrology - groundwater levels	Water levels (centimetres); duration of levels; hydraulic gradients; water supply levels	Maintain, or where necessary restore, appropriate natural hydrological regimes necessary to support the natural structure and functioning of the habitat	Fen habitats require high groundwater levels (i.e. water levels at or above the ground surface) for a large proportion of the calendar year (i.e. duration of mean groundwater level). Fen groundwater levels are controlled by regional groundwater levels in the contributing catchment area (which sustain the hydraulic gradients of the fen groundwater table). Regional abstraction of groundwater may affect fen groundwater levels
Ecosystem function: hydrology - surface water flow	Drain density and form	Maintain, or where necessary restore, as close as possible to natural or semi-natural, drainage conditions	Drainage, either within or surrounding the fen habitat, can result in the drawdown of the groundwater table. The depth, geometry and density of drainage (hydromorphology) will indicate the scale and impact on fen hydrology. Drainage can result in loss of characteristic species and transition to drier habitats
Ecosystem function: water quality	Various	Maintain appropriate water quality, particularly pH and nutrient levels, to support the natural structure and functioning of the habitat	Fens receive natural levels of nutrients (e.g. iron, magnesium and calcium) from water sources. However, they are generally poor in nitrogen and phosphorus, with the latter tending to be the limiting nutrient under natural conditions. Water supply should be also relatively calcium-rich
Vegetation composition: community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	The entire diversity of alkaline fen vegetation communities present in the SAC is currently unknown. Information on the vegetation communities associated with alkaline fens is provided by O'Neill et al. (in prep.). See also the Irish Vegetation Classification (Perrin, 2018; <a href="http://www.biodiversityireland.ie/projects/ivc-classification-explorer">www.biodiversityireland.ie/projects/ivc-classification-explorer</a> )

Vegetation composition: typical brown mosses	Percentage cover at a representative number of monitoring stops	Maintain adequate cover of typical brown moss species	For lists of typical bryophyte species, including high quality indicator species, see O'Neill et al. (in prep.). Species recorded at Lough Shesk and Newtown Lough include: <i>Calliergon giganteum</i> , <i>Scorpidium scorpioides</i> , <i>Campyllum stellatum</i> , <i>Bryum pseudotriquetrum</i> , <i>Fissidens adianthoides</i> , <i>Scorpidium scorpioides</i> , <i>Calliergonella cuspidata</i> and <i>Ctenidium molluscum</i> (NPWS internal files)
Vegetation composition: typical vascular plants	Percentage cover at a representative number of monitoring stops	Maintain adequate cover of typical vascular plant species	For lists of typical vascular plant species for the different vegetation communities, including high quality indicators, see O'Neill et al. (in prep.). Typical species recorded in the habitat in the SAC include black bog-rush ( <i>Schoenus nigricans</i> ), dioecious sedge ( <i>C. dioica</i> ) and common butterwort ( <i>Pinguicula vulgaris</i> ) (NPWS internal files)
Vegetation composition: native negative indicator species	Percentage cover at a representative number of monitoring stops	Cover of native negative indicator species at insignificant levels	Negative indicators include species not characteristic of the habitat and species indicative of undesirable activities such as overgrazing, undergrazing, nutrient enrichment, agricultural improvement or impacts on hydrology. Native negative indicators may include <i>Anthoxanthum odoratum</i> , <i>Epilobium hirsutum</i> , <i>Holcus lanatus</i> , <i>Juncus effusus</i> , <i>Phragmites australis</i> and <i>Ranunculus repens</i> . See O'Neill et al. (in prep.)
Vegetation composition: non-native species	Percentage cover at a representative number of monitoring stops	Cover of non-native species less than 1%	Attribute and target based on O'Neill et al. (in prep.). Non-native species can be invasive and have deleterious effects on native vegetation. A low target is set as non-native species can spread rapidly and are most easily dealt with when still at lower abundances
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on O'Neill et al. (in prep.). Scrub and trees will tend to invade if fen conditions become drier
Vegetation composition: algal cover	Percentage cover at, and in local vicinity of, a representative number of monitoring stops	Cover of algae less than 2%	Attribute and target based on O'Neill et al. (in prep.). Algal cover is indicative of nutrient enrichment from multiple sources (McBride et al., 2011)
Vegetation structure: vegetation height	Percentage cover at a representative number of monitoring stops	At least 50% of the live leaves/flowering shoots are more than either 5cm or 15cm above ground surface depending on community type	Attribute and target based on O'Neill et al. (in prep.). While grazing may be appropriate in this habitat, excessive grazing can reduce the ability of plant species to regenerate reproductively and maintain species diversity, especially if flowering shoots are cropped during the growing season
Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of monitoring stops	Cover of disturbed bare ground not more than 10%	Attribute and target based on O'Neill et al. (in prep.). While grazing may be appropriate in this habitat, excessive areas of disturbed bare ground may develop due to unsuitable grazing regimes. Disturbance can include hoof marks, wallows, human footprints, vehicle and machinery tracks. Excessive disturbance can result in loss of characteristic species and presage erosion for peatlands
Physical structure: tufa formations	Percentage cover in local vicinity of a representative number of monitoring stops	Disturbed proportion of vegetation cover where tufa is present is less than 1%	Attribute and target based on O'Neill et al. (in prep.)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat; maintain features of local distinctiveness, subject to natural processes	This includes species on the Flora (Protection) Order, 2015 and/or Red Lists (Byrne et al., 2009; Regan et al., 2010; Lockhart et al., 2012; Wyse Jackson et al., 2016, etc.). The Near Threatened species (Wyse Jackson et al., 2016) round-leaved wintergreen ( <i>Pyrola rotundifolia</i> ) has been recorded in the habitat around Newtown Lough in the SAC (NPWS internal files)
Transitional areas between fen and adjacent habitats	Hectares; distribution	Maintain adequate transitional areas to support/protect the alkaline fen ecosystem and the services it provides	In many cases, fens transition to other wetland habitats. It is important that the transitional areas between fens and other habitats are maintained in as natural condition as possible in order to protect the functioning of the fen

## Conservation Objectives for : River Boyne and River Blackwater SAC [002299]

### 91E0 Alluvial forests with *Alnus glutinosa* and *Fraxinus excelsior* (Alno-Padion, Alnion incanae, Salicion albae)\*

To restore the favourable conservation condition of Alluvial forests with *Alnus glutinosa* and *Fraxinus excelsior* (Alno-Padion, Alnion incanae, Salicion albae)\* in River Boyne and River Blackwater SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes. See map 3 for surveyed woodland areas	Alluvial forests with <i>Alnus glutinosa</i> and <i>Fraxinus excelsior</i> (Alno-Padion, Alnion incanae, Salicion albae)* is present within River Boyne and River Blackwater SAC. As part of the National Survey of Native Woodlands (NSNW), the sub-sites Grove Island (NSNW site code 688) and Yellow Island (752) were surveyed by Perrin et al. (2008). Yellow Island (code 752) was also included in national monitoring surveys (O'Neill and Barron, 2013; Daly et al., in prep.). Map 3 shows the minimum area of alluvial forests within the SAC, which is estimated to be 16.7ha (Perrin et al., 2008; Daly et al., in prep.). It is important to note that further unsurveyed areas may be present within the SAC
Habitat distribution	Occurrence	No decline, subject to natural processes. The surveyed woodland locations are shown on map 3	Distribution based on Perrin et al. (2008) and Daly et al. (in prep.). It is important to note that further unsurveyed areas may be present within the SAC
Woodland size	Hectares	Area stable or increasing. Where topographically possible, "large" woods at least 25ha in size and "small" woods at least 3ha in size	The target areas for individual woodlands aim to reduce habitat fragmentation and benefit those species requiring 'deep' woodland conditions (Peterken, 2002). In some cases, topographical constraints may restrict expansion
Woodland structure: cover and height	Percentage; metres; centimetres	Total canopy cover at least 30%; median canopy height at least 7m; native shrub layer cover 10-75%; native herb/dwarf shrub layer cover at least 20% and height at least 20cm; bryophyte cover at least 4%	The target aims for a diverse structure with a canopy containing mature trees, shrub layer with semi-mature trees and shrubs, and well-developed field layer (herbs, graminoids and dwarf shrubs) and ground layer (bryophytes). Assessment criteria are described in Daly et al. (in prep.) and O'Neill and Barron (2013)
Woodland structure: community diversity and extent	Hectares	Maintain diversity and extent of community types	The Boyne River Islands are an example of gallery forests of willows ( <i>Salicion albae</i> ), which occur alongside river channels and on river islands, where tree roots are almost continuously submerged (Daly et al., in prep.). Grove Island (NSNW site code 688) and Yellow Island (752) were assigned by Perrin et al. (2008) to the <i>Salix triandra – Urtica dioica</i> vegetation type (2h) of the <i>Fraxinus excelsior – Hedera helix</i> group. This corresponds to the <i>Salix fragilis – Calystegia sepium</i> sub-community (WL3Di) of the Irish Vegetation Classification (Perrin, 2016; www.biodiversityireland.ie/projects/ivc-classification-explorer)
Woodland structure: natural regeneration	Seedling: sapling: pole ratio	Seedlings, saplings and pole age-classes of target species for 91E0* woodlands and other native tree species occur in adequate proportions to ensure survival of woodland canopy	The target species for 91E0* are alder ( <i>Alnus glutinosa</i> ), ash ( <i>Fraxinus excelsior</i> ) and willows ( <i>Salix</i> spp.). Assessment criteria are described in Daly et al. (in prep.) and O'Neill and Barron (2013)



Hydrological regime: flooding depth/height of water table	Metres	Appropriate hydrological regime necessary for maintenance of alluvial vegetation	Periodic flooding is essential to maintain alluvial woodlands along river and lake floodplains, but not for woodland around springs/seepage areas. Much of the river channel within the SAC was subject to arterial drainage schemes. Natural flood-plains now exist along only limited stretches of river (NPWS internal files)
Woodland structure: dead wood	Number per hectare	At least 19 stems/ha of dead wood of at least 20cm diameter	Dead wood is a valuable resource and an integral part of a healthy, functioning woodland ecosystem
Woodland structure: veteran trees	Number per hectare	No decline	Veteran trees are important habitats for bryophytes, lichens, saproxylic organisms and some bird species. Their retention is important to ensure continuity of habitats/niches and propagule sources
Woodland structure: indicators of local distinctiveness	Occurrence; population size	No decline in distribution and, in the case of red listed and other rare or localised species, population size	Includes ancient or long-established woodlands (see Perrin and Daly, 2010), archaeological and geological features as well as red listed and other rare or localised species
Woodland structure: indicators of overgrazing	Occurrence	All five indicators of overgrazing absent	There are five indicators of overgrazing within 91E0*: topiary effect on shrubs and young trees, browse line on mature trees, abundant dung, severe recent bark stripping, and trampling (Daly et al., in prep.)
Vegetation composition: native tree cover	Percentage	No decline. Native tree cover at least 90% of canopy; target species cover at least 50% of canopy	The target species for 91E0* are alder ( <i>Alnus glutinosa</i> ), ash ( <i>Fraxinus excelsior</i> ) and willows ( <i>Salix</i> spp.) (Daly et al., in prep.; O'Neill and Barron, 2013)
Vegetation composition: typical species	Occurrence	At least 1 target species for 91E0* woodlands present; at least 6 positive indicator species for 91E0* woodlands present	A variety of typical native species should be present, depending on woodland type. The target species for 91E0* are alder ( <i>Alnus glutinosa</i> ), ash ( <i>Fraxinus excelsior</i> ) and willows ( <i>Salix</i> spp.). Positive indicator species for 91E0* are listed in Daly et al. (in prep.) and O'Neill and Barron (2013)
Vegetation composition: negative indicator species	Occurrence	Negative indicator species cover not greater than 10%; regeneration of negative indicator species absent	Negative indicator species (i.e. any non-native species, including herbaceous species) should be absent or under control. The canopy at Grove Island (NSNW site code 688) and Yellow Island (752) is dominated by a range of <i>Salix</i> spp. ( <i>S. cinerea</i> , <i>S. triandra</i> , <i>S. fragilis</i> , <i>S. viminalis</i> ) (Perrin et al., 2008). Although the latter two are not native to Ireland, an exception is made for these species where they occur within gallery woodland (Daly et al., in prep.). Perrin et al. (2008) recorded some sycamore ( <i>Acer pseudoplatanus</i> ) in the canopy at Grove Island (NSNW site code 688). Daly et al. (in prep.) found that the recent arrival of the invasive non-native herb Himalayan balsam ( <i>Impatiens glandulifera</i> ) at Yellow Island (752) has caused significant negative impacts to the alluvial forest habitat
Vegetation composition: problematic native species	Percentage	Cover of common nettle ( <i>Urtica dioica</i> ) less than 75%	Common nettle ( <i>Urtica dioica</i> ) is a positive indicator species for 91E0* but, in some cases, it may become excessively dominant. Increased light and nutrient enrichment are factors which favour proliferation of common nettle (Daly et al., in prep.)

## Conservation Objectives for : River Boyne and River Blackwater SAC [002299]

### 1099 River Lamprey *Lampetra fluviatilis*

To restore the favourable conservation condition of River Lamprey (*Lampetra fluviatilis*) in River Boyne and River Blackwater SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Percentage of river accessible	Restore access to all water courses down to first order streams	Artificial barriers can block or impede the passage of upstream migrating lamprey, thereby restricting access to spawning areas (Gargan et al., 2011; Rooney et al., 2015). There are a number of weirs along the lower sections of the Boyne main channel, the most substantial of these are located at Slane and downstream of Navan at Blackcastle. Efforts to trap adult river lamprey were undertaken at four locations throughout the catchment during November 2014 to April 2015. This was augmented in April 2015 by an extensive fyke-netting survey (n=26 sites). No adult river lamprey were encountered, with the only record to date being a dead individual from the River Boyne at Slane in late March 2015 (Gallagher et al., 2016). On the Boyne main channel, there is ideal spawning habitat both upstream and downstream of the weir at Blackcastle but spawning has not been observed at these locations to date
Distribution of larvae	Number of positive sites in 2nd order channels (and greater), downstream of spawning areas	Not less than 50% of sample sites with suitable habitat positive for larval brook/river lamprey	It is not possible to distinguish between larval brook and river lamprey in the field and they are therefore considered together in assessing conservation status. A survey of the Boyne catchment in 2015 recorded n=583 <i>Lampetra</i> spp. larvae from n=102 sites (Gallagher et al., 2016). As stated, the weirs in the lower main stem are a significant impediment to river lamprey passage and, for this reason, these larvae are considered to be mainly, if not all, brook lamprey. To achieve favourable condition <i>Lampetra</i> spp. should, as a minimum, be present in not less than 50% of all sampling sites surveyed with suitable habitat present within the natural range (JNCC, 2015). <i>Lampetra</i> spp. larvae were recorded from 72% of sites indicating a pass for this target. Distribution remained similar to a 2005 survey (O'Connor, 2006) although larvae continued to be absent from the Boycetown and Skane Rivers, as well as the upper reaches of the Kells Blackwater system
Population structure of larvae	Number of age/size classes	At least three age/size classes of larval brook/river lamprey present	The target of at least three age/size classes is based on guidance from JNCC (2015). Larvae typically range from 10-150mm in length and this corresponds to up to six age classes. A broad range of size classes (12-153mm), including young-of-year larvae, was recorded from the 2015 Boyne catchment-wide survey indicating a pass for this target. However, given the issue of artificial barriers on the River Boyne, it is likely that this value pertains to brook lamprey, as previously stated
Larval lamprey density in fine sediment	Larval lamprey/m <sup>2</sup>	Mean density of brook/river larval lamprey in sites with suitable habitat more than 5/m <sup>2</sup>	A target mean density of more than 5/m <sup>2</sup> larvae in sites with suitable habitat is required to achieve favourable condition (JNCC, 2015). In the Boyne survey a mean density of 6/m <sup>2</sup> <i>Lampetra</i> spp. larvae (n=583) was obtained. A number of tributaries did not achieve a pass for this target, including the Athboy/Tremblestown, Boycetown, Deel, Skane and Stonyford Rivers. Again, the overall mean density value is most likely indicative of the status of brook lamprey in the Boyne catchment

Extent and distribution of spawning nursery habitat	m <sup>2</sup> and occurrence	No decline in extent and distribution of spawning and nursery beds	<p>This target is based on spawning and nursery bed mapping during targeted larval lamprey monitoring surveys. River lamprey spawn in clean gravels in flowing water where they excavate shallow nests. While coarse substrate is required for spawning, the close proximity of nursery areas comprising mainly sand/silt are necessary for the development of larvae. The 2015 Boyne survey recorded adequate spawning and nursery habitat availability within the catchment (Gallagher et al., 2016). However, the sequence of weirs in the lower main channel of the Boyne represents a significant impediment to upstream passage. In addition, this lower section of river is in a degraded hydromorphological state with impounding and, therefore, poor habitat availability for spawning</p>
---	-------------------------------	--	--

---

## Conservation Objectives for : River Boyne and River Blackwater SAC [002299]

### 1106 Salmon *Salmo salar*

To restore the favourable conservation condition of Atlantic Salmon (*Salmo salar*) in River Boyne and River Blackwater SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution: extent of anadromy	Percentage of river accessible	100% of river channels down to second order accessible from estuary	Artificial barriers block salmon's upstream migration, thereby limiting species to lower stretches and restricting access to spawning areas. There are multiple barriers to fish migration in the Boyne system
Adult spawning fish	Number	Conservation limit (CL) for each system consistently exceeded	A conservation limit (CL) is defined by the North Atlantic Salmon Conservation Organisation (NASCO) as "the spawning stock level that produces long-term average maximum sustainable yield as derived from the adult to adult stock and recruitment relationship". The target is based on the Technical Expert Group on Salmon's (TEGOS) annual model output of CL attainment levels. See Gargan et al. (2021) for further details. Stock estimates are either derived from direct counts of adults (rod catch, fish counter) or indirectly by fry abundance counts. The Boyne is significantly below its CL
Salmon fry abundance	Number of fry/5 minutes electrofishing	Maintain or exceed 0+ fry mean catchment-wide abundance threshold value. Currently set at 17 salmon fry/5 minutes sampling	Target is threshold value for rivers currently exceeding their conservation limit (CL)
Out-migrating smolt abundance	Number	No significant decline	Smolt abundance can be negatively affected by a number of impacts such as estuarine pollution, predation and sea lice ( <i>Lepeophtheirus salmonis</i> )
Number and distribution of redds	Number and occurrence	No decline in number and distribution of spawning redds due to anthropogenic causes	Salmon spawn in clean gravels. There is restricted habitat for salmon in the Boyne and habitat rehabilitation programmes have been undertaken in sections of the catchment
Water quality	EPA Q value	At least Q4 at all sites sampled by EPA	Q values based on triennial water quality surveys carried out by the Environmental Protection Agency (EPA)

## Conservation Objectives for : River Boyne and River Blackwater SAC [002299]


### 1355 Otter *Lutra lutra*


To maintain the favourable conservation condition of Otter (*Lutra lutra*) in River Boyne and River Blackwater SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Distribution	Percentage positive survey sites	No significant decline	Measure based on standard otter survey technique. Favourable Conservation Status (FCS) target, based on 1980/81 survey findings, is 88% in SACs. Current range is estimated at 93.6% (Reid et al., 2013)
Extent of terrestrial habitat	Hectares	No significant decline. Area mapped and calculated as 447.6ha along river banks/ lake shoreline/around ponds	No field survey. Areas mapped to include 10m terrestrial buffer, identified as critical for otters (NPWS, 2007), along rivers and around water bodies
Extent of freshwater (river) habitat	Kilometres	No significant decline. Length mapped and calculated as 263.3km	No field survey. River length calculated on the basis that otters will utilise freshwater habitats from estuary to headwaters (Chapman and Chapman, 1982)
Extent of freshwater (lake) habitat	Hectares	No significant decline. Area mapped and calculated as 31.6ha	No field survey. Area mapped based on evidence that otters tend to forage within 80m of the shoreline (NPWS, 2007)
Couching sites and holts	Number	No significant decline	Otters need lying up areas throughout their territory where they are secure from disturbance (Kruuk and Moorhouse, 1991; Kruuk, 2006)
Fish biomass available	Kilograms	No significant decline	Broad diet that varies locally and seasonally, but dominated by fish, in particular salmonids, eels and sticklebacks in freshwater (Bailey and Rochford, 2006; Reid et al., 2013)
Barriers to connectivity	Number	No significant increase	Otters will regularly commute across stretches of open water up to 500m, e.g. between the mainland and an island; between two islands; across an estuary (De Jongh and O'Neill, 2010). It is important that such commuting routes are not obstructed



**Legend**

 River Boyne and River Blackwater SAC 002299




 **An Roinn Tithíochta, Rialtais Áitiúil agus Oidhreacht**  
 Department of Housing, Local Government and Heritage

**MAP 1:  
 RIVER BOYNE AND RIVER BLACKWATER SAC  
 CONSERVATION OBJECTIVES  
 SAC DESIGNATION**

Map to be read in conjunction with the NPWS Conservation Objectives Document

**SITE CODE:  
 SAC 002299; version 3.02.  
 CO. CAVAN / LOUGH / MEATH / WESTMEATH**


0 2 4 8 Kilometres




The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.  
 Ordnance Survey of Ireland Licence No OSI-NMA-014. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbheithnithe a déanamh ar theorainneacha na gceantar comharthaíthe. Suirbhéarachta Ordoanáis na hÉireann Ceadúnas Uimh OSI-NMA-014. © Suirbhéarachta Ordoanáis na hÉireann Rialtas na hÉireann

**N**




**Date: January 2021**



**Legend**

- River Boyne and River Blackwater SAC 002299
- River Boyne and River Blackwater SPA 004232
- Boyne Coast and Estuary SAC 001957
- Boyne Estuary SPA 004080

Ordnance Survey Ireland 2018.



**Legend**

- 91E0 \*Alluvial forests with *Alnus glutinosa* and *Fraxinus excelsior* (*Alno-padion*, *Alnion incanae*, *Salicion albae*)
- WN6 Wet willow-alder-ash woodland
- River Boyne and River Blackwater SAC 002299